

## Durham Research Online

---

### Deposited in DRO:

21 October 2015

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Streuff, K. and Forwick, M. and Szczuciński, W. and Andreassen, K. and O'Cofaigh, C. (2015) 'Submarine landform assemblages and sedimentary processes related to glacier surging in Kongsfjorden, Svalbard.', *Arktos.*, 1 . p. 14.

### Further information on publisher's website:

<http://dx.doi.org/10.1007/s41063-015-0003-y>

### Publisher's copyright statement:

© The Author(s) 2015. This article is published with open access at Springerlink.com Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

### Additional information:

## Use policy

---

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

# arktos - The Journal of Arctic Geosciences

## Submarine landform assemblages and sedimentary processes related to glacier surging in Kongsfjorden, Svalbard --Manuscript Draft--

<b>Manuscript Number:</b>																
<b>Full Title:</b>	Submarine landform assemblages and sedimentary processes related to glacier surging in Kongsfjorden, Svalbard															
<b>Article Type:</b>	Original Article															
<b>Funding Information:</b>	<table> <tr> <td>Norges Forskningsråd</td> <td>Karin Andreassen</td> </tr> <tr> <td>Statoil</td> <td>Karin Andreassen</td> </tr> <tr> <td>BG Group</td> <td>Karin Andreassen</td> </tr> <tr> <td>Det Norske Oljeselskap ASA</td> <td>Karin Andreassen</td> </tr> <tr> <td>European Commission FP7-People 2012 Initial Training Networks "Glaciated North Atlantic Margins (GLANAM)"</td> <td>Miss Katharina Teresa Streuff</td> </tr> <tr> <td>Polish Ministry of Science and Higher Education (IP2010040970)</td> <td>Witold Szczucinski</td> </tr> <tr> <td>Polish National Science Center (2013/10/E/ST10/00166)</td> <td>Witold Szczucinski</td> </tr> </table>		Norges Forskningsråd	Karin Andreassen	Statoil	Karin Andreassen	BG Group	Karin Andreassen	Det Norske Oljeselskap ASA	Karin Andreassen	European Commission FP7-People 2012 Initial Training Networks "Glaciated North Atlantic Margins (GLANAM)"	Miss Katharina Teresa Streuff	Polish Ministry of Science and Higher Education (IP2010040970)	Witold Szczucinski	Polish National Science Center (2013/10/E/ST10/00166)	Witold Szczucinski
Norges Forskningsråd	Karin Andreassen															
Statoil	Karin Andreassen															
BG Group	Karin Andreassen															
Det Norske Oljeselskap ASA	Karin Andreassen															
European Commission FP7-People 2012 Initial Training Networks "Glaciated North Atlantic Margins (GLANAM)"	Miss Katharina Teresa Streuff															
Polish Ministry of Science and Higher Education (IP2010040970)	Witold Szczucinski															
Polish National Science Center (2013/10/E/ST10/00166)	Witold Szczucinski															
<b>Abstract:</b>	<p>New high-resolution swath bathymetry data from inner Kongsfjorden, Svalbard, reveal characteristic landform assemblages formed during and after surges of tidewater glaciers, and provide new insights into the dynamics of surging glaciers. Glacier front oscillations and overriding related to surge activity lead to the formation of overridden moraines, glacial lineations of two types, terminal moraines, associated debris lobes and De Geer moraines. In contrast to submarine landform assemblages from other Svalbard fjords, the occurrence of two kinds of glacial lineations and the presence of De Geer moraines suggest variability in the landforms produced by surge-type tidewater glaciers. All the landforms in inner Kongsfjorden were deposited during the last c. 150 years. Lithological and acoustic data from the innermost fjord reveal that suspension settling from meltwater plumes as well as ice rafting are dominant sedimentary processes in the fjord, leading to the deposition of stratified glacimarine muds with variable numbers of clasts. Reworking of sediments by glacier surging results in the deposition of sediment lobes containing massive glacimarine muds. Two sediment cores reveal minimum sediment accumulation rates related to the Kongsvegen surge from 1948; these were 30 cm a<sup>-1</sup> approximately 2.5 km beyond the glacier front shortly after surge termination, and rapidly dropped to an average rate of 1.8 cm a<sup>-1</sup> in ~1950, during glacier retreat.</p>															
<b>Corresponding Author:</b>	Katharina Teresa Streuff, MSc Durham University Durham, UNITED KINGDOM															
<b>Corresponding Author Secondary Information:</b>																
<b>Corresponding Author's Institution:</b>	Durham University															
<b>Corresponding Author's Secondary Institution:</b>																
<b>First Author:</b>	Katharina Teresa Streuff, MSc															
<b>First Author Secondary Information:</b>																
<b>Order of Authors:</b>	<table> <tr> <td>Katharina Teresa Streuff, MSc</td> </tr> <tr> <td>Matthias Forwick</td> </tr> <tr> <td>Witold Szczucinski</td> </tr> </table>		Katharina Teresa Streuff, MSc	Matthias Forwick	Witold Szczucinski											
Katharina Teresa Streuff, MSc																
Matthias Forwick																
Witold Szczucinski																

	Karin Andreassen
	Colm O'Cofaigh
<b>Order of Authors Secondary Information:</b>	
<b>Author Comments:</b>	
<b>Suggested Reviewers:</b>	<p>Kelly Hogan Marine geophysicist, British Antarctic Survey kelgan@bas.ac.uk Kelly works with geophysical data including subbottom profiler data and bathymetry and she also has a very good knowledge of sediment cores and the depositional processes in front of glaciers/ice streams. She investigated a similar area from Greenland.</p> <p>John Howe Head of the Biogeochemistry and Earth Science Department, Scottish Association for Marine Science john.howe@sams.ac.uk John is familiar with the area, having worked there himself in 2003. He knows the seafloor adjacent to our study area very well. He works with multibeam data and sediment cores himself and is thus very knowledgeable.</p> <p>Paul Dunlop Senior Lecturer, University of Ulster p.dunlop@ulster.ac.uk Paul is specialised in geomorphology and landforms related to glacial activity. He is very familiar with the sedimentary processes involved and has a good knowledge of marine geophysical data.</p> <p>Lilja R Bjarnadottir Project Leader MAREANO, Geological Survey of Norway lilja.bjarnadottir@ngu.no Lilja is the project leader of the MAREANO project which maps the ocean floor around Norway. She is thus extremely knowledgeable in the work with bathymetry data. She previously worked on several projects from the Barents Sea and is also very familiar with glacial landform assemblages.</p> <p>Astrid Lysa Geological Survey of Norway astrid.lysa@ngu.no Astrid is very familiar with the dynamics of glacial advances and retreats and has worked on Svalbard for a number of occasions. She is thus very familiar with the area and the topic.</p>

Noname manuscript No.  
(will be inserted by the editor)

# Submarine landform assemblages and sedimentary processes related to glacier surging in Kongsfjorden, Svalbard

Katharina Streuff · Matthias Forwick ·  
Witold Szczuciński · Karin Andreassen ·  
Colm Ó Cofaigh

Received: date / Accepted: date

**Abstract** New high-resolution swath bathymetry data from inner Kongsfjorden, Svalbard, reveal characteristic landform assemblages formed during and after surges of tidewater glaciers, and provide new insights into the dynamics of surging glaciers. Glacier front oscillations and overriding related to surge activity lead to the formation of overridden moraines, glacial lineations of two types, terminal moraines, associated debris lobes and De Geer moraines. In contrast to submarine landform assemblages from other Svalbard fjords, the occurrence of two kinds of glacial lineations and the presence of De Geer moraines suggest variability in the landforms produced by surge-type tidewater glaciers. All the landforms in inner Kongsfjorden were deposited during the last c. 150 years. Lithological and acoustic data from the innermost fjord reveal that suspension settling from meltwater plumes as well as ice rafting

Katharina Streuff:  
Department of Geology, UiT - The Arctic University of Norway, Postboks 6050 Langnes, N-9037 Tromsø, Norway, Tel.: +44 (0) 191 33 41943, Fax: +44 (0) 191 33 41801  
E-mail: [katharina.streuff@durham.ac.uk](mailto:katharina.streuff@durham.ac.uk)  
*Present address:* Department of Geography, Durham University, Science Site, South Road, Durham, DH13LE, United Kingdom

Matthias Forwick  
Department of Geology, UiT - The Arctic University of Norway, Postboks 6050 Langnes, N-9037 Tromsø, Norway

Witold Szczuciński  
Institute of Geology, Adam Mickiewicz University in Poznań, Maków Polnych 16, 61-606 Poznań, Poland

Karin Andreassen  
Department of Geology, UiT - The Arctic University of Norway, Postboks 6050 Langnes, N-9037 Tromsø, Norway  
*also at:* CAGE -Centre for Arctic Gas Hydrate, Environment and Climate, Department of Geology, UiT - The Arctic University of Norway, Tromsø, Norway

Colm Ó Cofaigh  
Department of Geography, Durham University, Science Site, South Road, Durham, DH13LE, United Kingdom



are dominant sedimentary processes in the fjord, leading to the deposition of stratified glacialine muds with variable numbers of clasts. Reworking of sediments by glacier surging results in the deposition of sediment lobes containing massive glacialine muds. Two sediment cores reveal minimum sediment accumulation rates related to the Kongsvegen surge from 1948; these were 30 cm a<sup>-1</sup> approximately 2.5 km beyond the glacier front shortly after surge termination, and rapidly dropped to an average rate of 1.8 cm a<sup>-1</sup> in ~1950, during glacier retreat.

**Keywords** submarine landforms · glacier surges · tidewater glaciers · multibeam bathymetry · lithology · Svalbard

## 1 Introduction

Glacier surges are cyclic switches between active and passive phases, during which the ice front may either advance rapidly (active), stagnate (transition), or retreat slowly (passive/quiescent phase; e.g. Meier and Post, 1969; Sharp, 1985; Raymond, 1987; Dowdeswell et al, 1995; Gilbert et al, 2002). They generally occur independently of climate and are triggered internally through, for example, changes in glacier hydrology or basal thermal regime (Meier and Post, 1969; Kamb, 1987; Raymond, 1987; Sharp, 1988). Surges are common on Svalbard, where they have a generally longer duration than elsewhere, with active phases of between 4 and 10 years and quiescent phases of between 50 and 500 years (Dowdeswell et al, 1991; Murray et al, 1998; Benn and Evans, 2010). Many Svalbard glaciers have been identified as surge-type (e.g. Liestøl, 1969; Dowdeswell et al, 1991; Hagen, 1993; Plassen et al, 2004; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008), with surges well-documented from the past c. 180 years, but only two older examples (Paulabreen and Nathorstbreen; Liestøl, 1969; Hagen, 1993; Hald et al, 2001; Kristensen et al, 2009; Kempf et al, 2013).

Glacier surges lead to the formation of characteristic landform assemblages, which are revealed when the glacier retreats (e.g. Sharp, 1985; Solheim, 1991; Boulton et al, 1996; Evans and Rea, 1999; Evans et al, 1999; Evans, 2003; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008). Landforms related to glacier surges in submarine settings have been described from several Spitsbergen fjords (Boulton et al, 1996; Plassen et al, 2004; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008; Baeten et al, 2010; Flink et al, 2015). The landform models suggested thus far include overridden recessional moraines, (mega-scale) glacial lineations, terminal moraines with associated sediment lobes on their distal slopes, eskers, annual push moraines and crevasse-squeeze ridges, the latter suggested to be the only feature diagnostic of a glacier surge (Sharp, 1985; Ottesen et al, 2008). The high detail preserved in submarine environments offers invaluable insights into the processes controlling landform genesis, and, together with lithological records from sediment cores, enables a better understanding of tidewater glacier sedimentation and dynamics (e.g. Boulton et al, 1996; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008).

In this paper we present acoustic data (swath bathymetry and high-resolution seismic data) and lithological analyses of two sediment cores from inner Kongsfjorden, Svalbard. We describe and interpret submarine landform assemblages and deposits related to glacier surges and show that such assemblages are more diverse than previously suggested.

## 2 Study area

Kongsfjorden is located on northwestern Spitsbergen, the largest island of the Svalbard archipelago. It is the southern branch of the Kongsfjorden-Krossfjorden fjord system (78°50'N, 11°40'E, and 79°04'N, 12°40'E; Fig. 1). Kongsfjorden and Krossfjorden merge towards the open sea, where a large submarine trough, Kongsfjordrenna, channelled fast-flowing ice streams during the last glacial (e.g. Ottesen et al, 2005; Ingólfsson and Landvik, 2013). Kongsfjorden is approximately 20 km long and between 4 and 10 km wide. It covers an area of  $\sim 210 \text{ km}^2$ , and has a volume of  $29.4 \text{ km}^3$  (Ito and Kudoh, 1997). Water depths range from 350 m in the outer and central parts to  $<100 \text{ m}$  in the inner fjord. A detailed review on Kongsfjorden's climate and oceanography was provided by Svendsen et al (2002).

Seven tidewater glaciers terminate in Kongsfjorden: Løvlandbreen and Svansbreen form one tidewater front with Blomstrandbreen in the north of the fjord (Fig. 1) and will be summarized by the term "Blomstrandbreen" throughout this paper. Conwaybreen, Kongsbreen and Kronebreen dominate the east of the fjord with Kongsbreen terminating as two tidewater margins, one north and one south of Ossian Sarsfjellet (Fig. 1). Kongsvegen flows into the fjord from the south-east, adjacent to Kronebreen (Fig. 1). Three of these glaciers have been documented to be of surge-type, with Kronebreen and Kongsvegen experiencing respective surges in 1869 and 1948, and Blomstrandbreen surging in 1960 (Liestøl, 1988; Hagen, 1993).

## 3 Glacial history

The glacial history of the Svalbard archipelago during and since the Late Weichselian is documented to have occurred in three main stages: (a) ice advance, (b) full glaciation during the Last Glacial Maximum (LGM), and (c) ice retreat (see e.g. Elverhøi et al, 1995; Landvik et al, 1998; Jessen et al, 2010).

During initial advance, ice extended beyond the present coastline, reaching the shelf break between  $24,080 \pm 150$  and  $23,550 \pm 185 \text{ cal a BP}$  (calibrated years before present; Landvik et al, 1998; Jessen et al, 2010). Fast-flowing ice streams drained the Svalbard-Barents Sea Ice-Sheet via the main fjord systems on Svalbard, including Kongsfjorden (e.g. Ottesen et al, 2005, 2007; Ingólfsson and Landvik, 2013). A terminal moraine at the shelf break in southern Kongsfjordrenna was inferred to reflect maximum ice extent during the Late Weichselian (Ottesen et al, 2007). Deglaciation of the shelf and fjords on

west Spitsbergen began around  $20,500 \pm 500$  cal a BP and was interrupted by multiple glacier halts and/or re-advances (e.g. Ottesen et al, 2007; Baeten et al, 2010; Jessen et al, 2010; Forwick and Vorren, 2009, 2010; Kempf et al, 2013). The deglaciation of Kongsfjorden proper was documented as a two-stage recession initiated  $\sim 13,000$  cal a BP, leading to ice-free conditions by approximately 9,000 cal a BP (Lehman and Forman, 1992). Recent findings by Henriksen et al (2014), however, show that the ice stream in Kongsfjorden had retreated to the fjord mouth by 16,600 cal a BP, and that the deglaciation of the areas west of Blomstrandhalvøya was already complete before 14,400 ( $\pm 300$ ) cal a BP.

Asynchronous re-growth of Svalbard glaciers occurred after c. 9,000 cal a BP (e.g. Forwick and Vorren, 2007, 2009; Baeten et al, 2010; Forwick et al, 2010). Maximum Late Holocene glacier extents occurred either due to climatic cooling during the Little Ice Age or due to glacier surges (e.g. Liestøl, 1969; Dowdeswell et al, 1991; Hagen, 1993; Plassen et al, 2004; Ottesen and Dowdeswell, 2006; Mangerud and Landvik, 2007; Ottesen et al, 2008; Kempf et al, 2013).

## 4 Methods

Swath-bathymetry, sub-bottom profiler (chirp) data and sediment cores acquired in autumn 2010 with R/V Jan Mayen (now R/V Helmer Hanssen) from inner Kongsfjorden provide the basis for this study (Fig. 1). A Kongsberg Maritime Simrad EM 300 multibeam echo sounder was used to acquire the bathymetry data (max. resolution of 5 m). The instrument operated at a frequency of approximately 30 kHz and was calibrated using p-wave velocities for the water column obtained from CTD (conductivity, temperature, depth) measurements. The bathymetry data were supplemented with multibeam data from the Norwegian Hydrographic Survey gridded to a max. resolution of 5 m and visualised and interpreted using the Fledermaus v7 3D Visualization and Analyzing Software. Chirp data are exclusively available for the innermost part of Kongsfjorden (Fig. 1c), where profiles were recorded using a hull-mounted EdgeTech 3300-HM sub-bottom profiler operating at a pulse mode of 2-12 kHz and 3 ms, while the ping rate was set to 1.9 Hz. The profiles were processed in the EdgeTech Software and interpreted using SMT The Kingdom Suite.

Two gravity cores, 10JM-GlaciBar-GC01 (GC01) and 10JM-GlaciBar-GC02 (GC02), were retrieved with a 1900 kg heavy gravity corer with a 6 m long barrel. After retrieval the cores were divided into sections of up to 1 m in length and were subsequently stored at  $+4^{\circ}\text{C}$ . The core sites are located  $\sim 310$  m apart from each other, with GC01 ( $78^{\circ}55'50''\text{N}$ ,  $12^{\circ}20'49''\text{E}$ ; 50 m water depth; 286 cm length) recovered from the top of a sediment wedge (representing a debris lobe deposited from a glacier surge, see section 5 below), and GC02 ( $78^{\circ}55'59''\text{N}$ ,  $12^{\circ}20'36''\text{E}$ ; 53 m water depth; 339 cm length) from c. 130 m beyond this wedge (Fig. 1b). The p-wave velocity of the sediments was measured in 1 cm increments using a GEOTEK multi-sensor core logger (MSCL)

at UiT - The Arctic University of Norway prior to opening of the cores. Lithological logs are based on visual descriptions of the sediment surfaces as well as X-radiographs taken with a Philips Macrotank (5 mA; 80 kV; exposure times: 100 sec to 4 min). Sediment samples ( $\sim 1$  g every 10 cm) were measured with a Beckman Coulter LS13320 Laser Diffraction Particle Size Analyzer to obtain information on grain size distribution. Prior to measurements each sample was dissolved in 50 ml of water and homogenized in a shaker.

Sediment accumulation rates (SAR) for the last century were determined through complementary  $^{210}\text{Pb}$  and  $^{137}\text{Cs}$  analyses, as previously used for Svalbard fjords (Svendsen et al, 2002; Zajaczkowski et al, 2004; Zaborska et al, 2006; Szczuciński et al, 2009). Activities of both isotopes were measured with gamma spectroscopy using a Canberra GX2520 high-purity coaxial germanium detector at the Institute of Geology, Adam Mickiewicz University in Poznań, Poland. For this, sediment samples of about 20 g were taken from 10 cm thick intervals, dried and ground. Obtained activities were decay-corrected to the date of sampling, and the results are presented with a two-sigma standard deviation uncertainty range. From the decrease of excess  $^{210}\text{Pb}$  activities with sediment depth, SAR could be calculated (following McKee et al, 1983). Excess  $^{210}\text{Pb}$  activities were determined by taking the average supported activity from the sample below the region of radioactive decay, and subtracting it from the total activity. The independent SAR assessment was made using the first occurrence of  $^{137}\text{Cs}$  as a marker of the early 1950s (taken as 1952), its maximum activity peak as  $\sim 1962$  and the younger secondary activity peak as Chernobyl-related 1986 (e.g. Robbins and Edgington, 1975; Appleby, 2008). However, due to the possible loss of the core top sediment during the coring process, sediment mixing, variations in sediment accumulation rates and low activities of excess  $^{210}\text{Pb}$ , the calculated sediment accumulation rates should be treated as approximate values.

A digital terrain model (DTM) from 2009 with a vertical resolution of 5 m (Delmodell 5m 2009.13822\_33, courtesy of the Norwegian Polar Institute, provided on geodata.npolar.no) was supplemented with satellite imagery available on Google Earth (August 2015) and visualised in Esri ArcMap 10.2. Superficial crevasses were then mapped on all tidewater glaciers in Kongsfjorden within  $\sim 1$  km of the current glacier fronts.

## 5 Results and interpretations

### 5.1 Seafloor morphology

Landforms occurring in Kongsfjorden are described and interpreted in the following section. Their distribution is shown in Figure 2.

### 5.1.1 Large transverse ridges – overridden moraines

10 to 30 m high ridges are orientated generally transverse to the direction of ice flow and occur in front of Kronebreen/Kongsvegen and Blomstrandbreen (Fig. 2b). They are up to 300 m wide and around 1 km long. Their crests are round, smooth, cross-cut by streamlined bedforms (section 5.1.2) and are overprinted by small sharp-crested transverse ridges (section 5.1.4).

The large ridges are very similar to transverse ridges from Borebukta (Ottesen and Dowdeswell, 2006) and are thus interpreted to be moraines deposited by a tidewater glacier during an earlier phase of stagnation or retreat. These ridges were then overridden during a subsequent advance, leading to the formation of the streamlined bedforms described in section 5.1.2. The occurrence of several of these moraines in front of Kronebreen and Kongsvegen (Figs. 2b and 3a) suggests that here they represent recessional moraines deposited from repeated stillstands during overall retreat of the glacier. The single ridge in front of Blomstrandbreen (RB2, Fig. 2b), however, probably represents a terminal moraine (cf. section 5.1.3 below) from an earlier advance.

### 5.1.2 Streamlined bedforms – glacial lineations

Two types of streamlined bedforms are distinguished: (1) sets of parallel, 8 m high, 20-80 m wide, and up to 2 km long smooth-crested grooves and ridges, aligned parallel to the direction of ice flow (Figs. 2b, 3c) and (2) sets of parallel, ~2 m high, ~20 m wide and up to 600 m long sharp-crested grooves and ridges, also aligned parallel to the direction of ice flow and spaced at variable distances between 50 and 300 m (Figs. 2b, 3e). These latter features are confined to the areas in front of Kongsbreen North and Kronebreen/Kongsvegen, whereas the smoother groove-ridge features occur in front of Blomstrandbreen, Conwaybreen and Kongsbreen South. Similar features also appear in the outer fjord where they were likely formed during the last glacial (Fig. 2b; MacLachlan et al, 2010). With the exception of these latter features, all streamlined bedforms in Kongsfjorden are overprinted by small, transverse ridges (see section 5.1.4).

The two types of streamlined bedforms in Kongsfjorden are interpreted to be glacial lineations. Based on appearance and dimensions, the grooves and ridges are similar to (mega-scale) glacial lineations (MSGLs; e.g. Clark, 1994; Stokes and Clark, 2002; Ottesen et al, 2005; Ottesen and Dowdeswell, 2006; Andreassen et al, 2007; Baeten et al, 2010; Flink et al, 2015), which are closely associated with fast ice-flow (King et al, 2009). Even though lineations in Kongsfjorden are much shorter (max. 2 km) than MSGLs described from elsewhere (up to 70 km; Clark, 1993), the majority have elongation ratios of 10:1 or greater, technically classifying them as MSGLs. We infer that the Kongsfjorden lineations resulted from the same processes forming MSGLs, i.e. fast ice-flow (e.g. Stokes and Clark, 2002), when processes of erosion and re-deposition deform soft subglacial sediments into sets of grooves and ridges (cf. e.g. Tulaczyk et al, 2001; Ó Cofaigh et al, 2005; Ottesen et al, 2008; King

et al, 2009). In Kongsfjorden fast ice flow was probably due to the onset of the active phase of a glacier's surge-cycle. We note, however, that the differences in size and crest morphology between the two lineation types in Kongsfjorden indicates that their formation probably occurred under slightly different conditions. This issue is further discussed in section 6.1 below.

All glacial lineations in inner Kongsfjorden are inside the respective glaciers' maximum extents, so we interpret them to have formed within the last ~150 years (see section 6.2 below).

### *5.1.3 Large transverse ridges and lobe-shaped deposits – terminal moraines and debris lobes*

Seven large transverse ridges occur in Kongsfjorden, three in front of Blomstrandbreen (RB1, RB2 (overridden), RB3; numbered from distal to proximal, Fig. 2b,c), two in front of Conwaybreen and Kongsbreen North (RC1 and RC2) and three in front of Kronebreen and Kongsvegen (RK1, RK2 and RK3; Fig. 2c). They are 15 to 35 m high, between 500 and 2000 m long and several hundred meters wide (Fig. 4). Ridges in front of Blomstrandbreen and Kronebreen/Kongsvegen are separated into several segments by the surrounding islands (Fig. 2c). The ridges occur at distances of between 3 and 9 km from the present ice margins and are characterized by generally steeper proximal and gentler distal flanks (Fig. 4 d,e). RB3 differs from the other ridges by being symmetrical in cross-section and narrower (max. 100 m); it also has a much sharper crest (see Fig. 5d). With the exception of RB3 and RC1 the ridges occur in close association with lobe-shaped deposits on their distal flanks (Figs. 2c, 4b,c). These lobes occur as single deposits (up to 360 m wide and 600 m long) or in sets of (partly superimposed) tongue-shaped landforms. The latter can cover areas of up to 5 km<sup>2</sup>. The lobes typically occur at water depths between 15 and 50 m, but one lobe in the southwestern part of the fjord extends down to c. 110 m.

In front of the Kronebreen/Kongsvegen ice margin two lobes have very similar characteristics, but are dissociated from terminal moraines (Fig. 2c). They are separated by an approximately 3 m high elevation. Both features are located directly at the glacier margin and cover areas of 0.06 km<sup>2</sup> (380 x 160 m<sup>2</sup>) and 0.04 km<sup>2</sup> (500 x 85 m<sup>2</sup>), respectively. Chirp data reveal that these features are buried beneath stratified sediments (cf. section 5.2, Fig. 6c).

The large transverse ridges are inferred to be terminal moraines marking the maximum extent of, in most cases several, glacier advances. These ridges could be of glaciotectonic origin and reflect pushed-up, folded and/or thrust sub- or proglacial sediments, as described for other areas on Svalbard (e.g. Solheim and Pfirman, 1985; Boulton, 1986; Boulton et al, 1996; Plassen et al, 2004; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008; MacLachlan et al, 2010). We suggest that the lobe-shaped landforms are debris lobes that represent either (1) a product of downslope mass-transport of glacial sediment deposited from quasi-continuous slope failure on the distal side of the moraine during or after maximum ice extent, or (2) glacier-outwash fans formed by

meltwater-related processes during maximum extent of the glacier (e.g. Boulton et al, 1996; Plassen et al, 2004; Ottesen et al, 2008; Kristensen et al, 2009). The sedimentary lobes in front of Kronebreen/Kongsvegen were probably formed from continuous high sediment supply from meltwater streams (cf. Trusel et al, 2010; Kehrl et al, 2011) and may indicate that the Kronebreen/Kongsvegen margin experienced a prolonged still-stand close to its 2010 position.

#### 5.1.4 Small, predominantly transverse ridges – De Geer moraines

Numerous small ridges, that are 1 – 5 m high, several hundred meters long, and around 30 m wide (Figs. 2d, 5), are observed in Kongsfjorden. Although generally transverse and (sub-)parallel to each other, they have variable orientations and, in some cases, have a "saw-tooth" pattern in planform (Fig. 2e,f). Individual ridges cross-cut each other in places, are spaced at irregular intervals between 5 and 100 m and can exhibit branching. They occur in water depths down to 150 m and have mostly sharp, symmetrical crests (Fig. 5). Some ridges are longer and straighter than others, may extend across the entire width of the fjord and have slightly sinuous crests orientated exclusively perpendicular to the direction of ice flow (Fig. 5d). About 55 of these latter ridges occur in the fjord basin southeast of Blomstrandbreen, where they have been deposited between the outermost terminal moraine and the current ice front (Fig. 2d). 45 ridges or segments thereof extend between the outermost moraine and the current ice front of Conwaybreen, whereas about 30 ridges were deposited in the proximal basin of Kongsbreen South (Fig. 2d). In front of Kronebreen/Kongsvegen, most of the small ridges are around 1 m high and show weakly defined crests and frequent changes in orientation (Figs. 5c,g).

Based on their dimensions and morphology the small ridges are interpreted as De Geer moraines (e.g. De Geer, 1940; Zilliacus, 1989). De Geer moraines form subaquatically and typically occur as sets of transverse, parallel, irregularly spaced ridges that are around 3 m high, several hundred m long and up to 30 m wide (Zilliacus, 1989; Lundqvist, 2000). Two main mechanisms have been proposed for their formation: (1) The ridges are annual end moraines composed of subglacial sediment pushed up at the glacier grounding line (e.g. De Geer, 1940; Boulton, 1986; Sollid, 1989; Larsen et al, 1991; Blake, 2000); (2) they are the product of sediment squeezed into basal crevasses (e.g. Hoppe, 1957; Strömberg, 1965; Zilliacus, 1989; Beaudry and Prichonnet, 1991). In Svalbard, similar ridges have been interpreted as either annual push moraines, created in front of tidewater glaciers by pushing during small winter re-advances, or as crevasse-squeeze ridges (Sharp, 1985; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008; Flink et al, 2015). As glaciers are believed to be especially crevassed when in the active phase of a surge cycle, the crevasse-squeeze ridges have been suggested to be the only landform definitively indicative of surge activity (Sharp, 1985; Ottesen and Dowdeswell, 2006). Following the recommendation of Lundqvist (1981), we interpret the small ridges in Kongsfjorden as De Geer moraines. Although they could have been formed by either of

the two suggested mechanisms (see above), we favour formation related to crevasse-squeezing. This issue is further discussed in section 6.1 below.

## 5.2 Seismostratigraphy

Chirp data reveal four acoustic facies in inner Kongsfjorden (Fig. 6). Facies 4 is stratigraphically oldest and is characterized by (semi-)transparent, acoustically massive reflections. It occasionally crops out along steeper slopes (Fig. 6). The thickness of Facies 4 ranges from 1 – 10 ms (two-way travel time – TWT), which converts to about 1 – 8 m, respectively (using  $1500 \text{ m s}^{-1}$  for all facies, the p-wave velocity determined from MSCL measurements). These are minimum thicknesses, however, as the max. penetration of the echosounder signal is  $\sim 26$  ms TWT, i.e.  $\sim 20$  m. Facies 3 is acoustically similar to Facies 4 and shows transparent and massive reflections. However, Facies 3 appears thicker than Facies 4 (min. 10 – 20 ms or 8 – 15 m thick) and has a wedge-like shape (as indicated by brown polygons in Fig. 6). Facies 3 only occurs on slopes, specifically the distal sides of terminal moraines, and close to the Kronebreen/Kongsvegen ice margin. It appears thicker at the foot of the slopes, where it generally onlaps onto Facies 4. However, on the distal side of RK3 it onlaps onto Facies 2 (Fig. 6b,d). Facies 2 is acoustically stratified with laterally (semi-)continuous, opaque, parallel reflections of variable strength (Fig. 6b,d). The facies is bounded by a variably strong, opaque, semi-continuous reflection at its top, which is largely parallel to the seabed, and a weaker semi-continuous reflection at its base. Facies 2 is around 5 ms ( $\sim 4$  m) thick and shows a downlapping character in some areas in Kongsfjorden (Fig. 6b,d). Facies 1 is similar to Facies 2, with parallel, (semi-)continuous, opaque and parallel reflections. It is bounded by the seabed on top (Fig. 6). We thus infer Facies 1 to be youngest. The thickness of Facies 1 ranges from 1 ms ( $\sim 1$  m) in distal and steep areas to max. 5 ms ( $\sim 4$  m) in ice-proximal areas. Both Facies 1 and 2 are abundant in Kongsfjorden and are particularly common in bathymetric depressions where together, they can be up to 11 ms (8 m) thick.

The stratigraphic relationship between the facies varies locally. In ice-distal areas and away from the terminal moraines Facies 4 directly underlies Facies 2, which, in turn, directly underlies Facies 1 (Fig. 6b,d). In the ice-proximal area of Kronebreen/Kongsvegen, however, Facies 1 directly overlies Facies 4 (Fig. 6). The differing relationship of Facies 3 to the other facies in the fjord suggests variable timing of deposition. Generally Facies 3 overlies and is thus considered younger than Facies 4. In the case of RK3, however, Facies 3 onlaps onto Facies 2 (Fig. 6d), indicating that at this locality Facies 3 is also younger than Facies 2.

The acoustically massive appearance of Facies 4 (and 3) is suggested to reflect a mixed lithological composition. This, together with the facies' distribution within the fjord, leads us to interpret Facies 4 as acoustic basement. As this facies dominates on steeper slopes and in hummocky terrain (Fig. 6), and numerous glacier advances occurred, it is likely that the sediments are



either bedrock or glacial sediments (cf. Forwick and Vorren, 2010), the latter possibly representing glacial till (sensu Evans et al, 2006) or subglacial till (see also Elverhøi et al, 1983). From its onlapping character, its wedge-like shape and its close association with distal moraine slopes we infer that Facies 3 represents the debris lobes described in section 5.1.3. Facies 3 is acoustically similar to sediment wedges from other Spitsbergen fjords (e.g. Plassen et al, 2004; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008), thus supporting our interpretations. The acoustic stratification of Facies 1 and 2 reflects repeated changes in the physical properties of the sediments most probably related to changes in grain size in a glacier-proximal environment (e.g. Plassen et al, 2004; Forwick et al, 2010; Forwick and Vorren, 2010). The differences in thickness between Facies 1 and 2 are explained by varying durations of deposition. In the case of Kongsvegen, which surged in 1948 (Hagen, 1993), the uppermost sediments between the terminal surge moraine and the glacier front, i.e. Facies 1, accumulated over a maximum of 62 years (1948 to 2010), while the sediments of Facies 1 and Facies 2 beyond the surge moraine (RK3) accumulated over a much longer time period (>100 years since the last advance, see Fig. 6d). The variable thickness of Facies 1 is probably related to the distance from the ice margin: in ice-proximal areas sedimentation rates are usually high, whereas they decrease exponentially with increasing distance from the glacier margin (e.g. Elverhøi et al, 1983). We infer the bottom reflector of Facies 1 to represent the surge surface of 1948, and the bottom reflector of Facies 2 to represent the surge surface from 1869, when Kronebreen surged (cf. Elverhøi et al, 1983; Hagen, 1993). The increased thicknesses of Facies 1 and 2 in bathymetric depressions might indicate elevated sediment input from the surrounding slopes. Facies 1, 2 and 3 have been sampled in the two sediment cores and their origin is discussed further in the next section.

### 5.3 Lithology

#### 5.3.1 Sedimentology

A total of three lithological units are distinguished: Unit 3 is found at the base of GC02 (205–339 cm; Fig. 7a) and is inferred to be the oldest unit. It contains lighter and darker sharp-based layers of (reddish) brown clayey silt, that are a few millimeters to several centimeters thick. Scattered clasts occur throughout. Unit 2 is a matrix-supported, soft and water-rich diamict occurring at the base of GC01 (165 to 286 cm). It has a sharp upper boundary and contains abundant clasts distributed in a matrix of massive clayey silt. Unit 1 forms the uppermost, and therefore youngest unit in both sediment cores and extends from 0–165 cm in GC01 and from 0–205 cm in GC02 (Fig. 7a). Unit 1, like Unit 3, contains stratified clayey silt; however, the clast content is much higher. Clasts occur in concentrated layers or clusters or as randomly distributed limestones (Fig. 7a). The boundary between units 3 and 1 in core GC02 is transitional.

We suggest that units 3 and 1 were deposited in a glacial marine environment where sedimentation occurred from suspension settling (mud) and ice rafting (clasts). This is in accordance with similar findings from Elverhøi et al (1980, 1983). Ice rafting probably occurred mainly from icebergs, but some contribution by sea ice is likely. The stratification of units 3 and 1 is probably related to recurring changes in the sediment source with varying contributions of meltwater from Kronebreen and Kongsvegen. Such variations could be due to changes in the glaciers' hydrology or seasonal variations in the rate of sediment delivery and discharge (cf. e.g. Szczuciński and Zajaczkowski, 2012). Based on the high water content, the massive internal structure and the variable grain size, we interpret the sediments of Unit 2 as reworked glacial marine sediments.

Measurements of the  $^{137}\text{Cs}$  activity in the two sediment cores show a presence of  $^{137}\text{Cs}$  in the upper 110 cm of both cores, with maximum activity at approximately 85 cm. A secondary peak of  $^{137}\text{Cs}$  activity appears in the uppermost 15 cm of GC02 (Fig. 7b). The  $^{210}\text{Pb}$  measurements reveal excess  $^{210}\text{Pb}$  in the upper 150 cm of both cores. These activities show an irregular profile in both cores, but generally decrease with depth (Fig. 7b).

The results of the  $^{137}\text{Cs}$  activity measurements indicate that the upper 110 cm in both cores, i.e. large parts of Unit 1, have likely been deposited after ~1950. Results further indicate that a sediment depth of approximately 85 cm in both cores is attributed to an age of 1962 AD (Fig. 7b). We thus infer that Unit 1 was deposited after the Kongsvegen surge in 1948 and reflects an increasingly ice distal environment during glacier retreat. As Unit 1 directly overlies Unit 2 in GC01, we deduce that the reworked sediments of Unit 2 were deposited analogous to the debris flows described in section 5.1.3 and represent the sediment lobe associated with RK1 (Figs. 6, 8). The stratigraphic relationship from the chirp data suggests the debris lobe from the 1948 Kongsvegen surge to be younger than Unit 3 (see section 5.3.2). We suggest, therefore, that the base of Unit 3 was deposited during quiescent-phase conditions after the previous surge, and its uppermost parts during the active phase of the 1948 surge. The relatively low number of clasts in Unit 3 could be explained by either a reduction in ice rafting or by high suspension rainout which swamped the contribution from IRD. The latter is most likely, as the high calving intensity of Kronebreen ( $0.2247 \text{ km}^3 \text{ a}^{-1}$ ; Błaszczyk et al, 2009), the configuration of local ocean and wind currents and the warm water temperatures resulting from increased inflow of Atlantic water are at odds with a low IRD delivery to the core sites (cf. Svendsen et al, 2002; Howe et al, 2003; Jernas et al, 2013).

### 5.3.2 Correlation of seismo- and lithostratigraphy

We correlate seismic Facies 3 with lithological Unit 2 (reworked glacial marine sediment deposited from debris flows triggered by surge activity), seismic Facies 2 with lithological Unit 3 (glacial marine muds deposited during glacier advance), and seismic Facies 1 with lithological Unit 1 (glacial marine muds deposited during glacier retreat; Fig. 8). The hiatus from Facies 1 to Facies 4

in the proximal parts of Kongsvegen/Kronebreen (Fig. 6) indicates that Unit 3 (Facies 2) was likely eroded here, probably as a result of glacial advance during the Kongsvegen surge.

### 5.3.3 Sediment accumulation rates

The age-depth relationship in both cores shows that the majority of Unit 1 was deposited with a calculated average sediment accumulation rate (SAR) of  $1.8 \text{ cm a}^{-1}$  (Fig. 7b). The results of the  $^{137}\text{Cs}$  activity measurements also indicate that the upper 15 cm in GC02 have probably been deposited since the Chernobyl-accident in 1986, which suggests a low SAR of  $0.6 \text{ cm a}^{-1}$  for this portion of Unit 1. In fact,  $^{210}\text{Pb}$  activity measurements suggest non-steady sedimentation conditions and thus variable SARs due to the irregular profile of excess  $^{210}\text{Pb}$ . The measurements also show a minimum SAR for the uppermost 150 cm in both cores of  $\sim 1.5 \text{ cm a}^{-1}$ , which is in accordance with the  $^{137}\text{Cs}$  dating. Note, however, that all rates presented here are minimum SARs, because the cores were taken with a gravity corer and the uppermost portion of the sediment cover may have been lost during sampling. This could, for example, explain the especially low SAR for the upper 15 cm of Unit 1.

The overall decrease in excess  $^{210}\text{Pb}$  activity with depth in both cores (see Fig. 7b) is related to its radioactive decay with time. Despite the fact that generally excess  $^{210}\text{Pb}$  is measurable in sediments up to c. 100 years old (Koide et al, 1972), no excess  $^{210}\text{Pb}$  could be detected in sediments older than c. 60 years in both cores (Fig. 7b). This could be due to the presence of sediments older than 100 years that were reworked and redeposited, or due to very high SAR causing dilution of the excess  $^{210}\text{Pb}$ . The latter is particularly likely, because the top of the sediment lobe deposited during the Kongsvegen surge in 1948 was inferred to be located at 165 cm in core GC01, while a sediment depth of 110 cm dates to 1950 (according to the  $^{137}\text{Cs}$  dating), thus suggesting a SAR of  $\sim 30 \text{ cm a}^{-1}$  (i.e. one order of magnitude higher than the SAR after 1950). It should be noted that this is a minimum sediment accumulation rate as the exact time of deposition of the sediment lobe remains unknown. We ascribe this high accumulation rate to the proximity of the glacier front, which, based on the distance between terminal moraine and core site, was located approximately 2.5 km from the core site. According to Trusel et al (2010) and Kehrl et al (2011), recent SARs at the immediate fronts of Kongsvegen and Kronebreen are  $>1 \text{ m a}^{-1}$ . As exceptionally high discharge of turbid meltwater is common during and immediately after glacier surges (e.g. Elverhøi et al, 1983; Gilbert et al, 2002; Björnsson et al, 2003), a former SAR of about  $30 \text{ cm a}^{-1}$  at the core site seems reasonable. Such high accumulation rates are also in accordance with the low number of IRD in lithological Unit 3, which was linked to high input of fine-grained material from meltwater plumes, masking the contribution of IRD (see section 5.3.1). As the rate of sedimentation from suspension in glacialmarine settings usually decreases exponentially with distance (Elverhøi et al, 1983; Farrow et al, 1983; Syvitski, 1989; Cowan and Powell, 1991; Szczuciński and Zajaczkowski, 2012),

the relatively rapid decline of SARs later on may be related to rapid glacier retreat and the increasing distance between core sites and sediment source. This effect may have been enhanced by trapping of sediments within the basin between the terminal moraine and the retreating glacier front (cf. Kempf et al, 2013).

## 6 Discussion

### 6.1 Surge signatures in Kongsfjorden and comparison with other Spitsbergen fjords

Each tidewater glacier in Kongsfjorden formed (1) glacial lineations during an advancing phase of fast ice flow, (2) terminal moraines when reaching maximum ice extent, (3) associated debris lobes on the moraines' distal slopes, and (4) De Geer moraines after the termination of the advance. In front of Blomstrandbreen and Kronebreen/Kongsvegen, overridden moraines reflect a terminal moraine and recessional moraines, respectively, which were modified during a later re-advance.

The observed landform assemblage is generally consistent with those observed for terrestrial surge-type glaciers (e.g. Evans and Rea, 1999; Evans and Twigg, 2002) and is also similar to submarine landform assemblages described for surge-type glaciers in other Spitsbergen fjords. This shows that the depositional models proposed by Plassen et al (2004), Ottesen and Dowdeswell (2006), Ottesen et al (2008) and Flink et al (2015) are largely applicable also in Kongsfjorden. However, the occurrence of two different types of glacial lineations in the same fjord, as well as the presence of De Geer moraines appears to be unique to Kongsfjorden.

#### *Various types of glacial lineations*

Features similar to the smoother grooves and ridges from Kongsfjorden also occur in other Svalbard fjords, including Billefjorden (Baeten et al, 2010), Lomfjorden, and Ymerbukta (Streuff, in prep.). Conversely, glacial lineations from Borebukta (Ottesen and Dowdeswell, 2006), Van Keulenfjorden, Rinderbukta (Ottesen et al, 2008), and Tempelfjorden (Forwick et al, 2010; Flink et al, 2015), are more comparable to the sharp-crested lineations from Kongsfjorden. Thus the formation processes for the latter were probably similar to those for MSGL, even though the lineations in Kongsfjorden are of a much smaller scale. Possible explanations for this could be e.g. a less deformable substratum beneath the glacier, slower ice flow, shorter advances leaving insufficient time for the formation of larger features, or thinner glacier ice. Such differences between the individual glaciers could also account for the presence of both types of glacial lineations in Kongsfjorden.

#### *De Geer moraines*

De Geer moraines have been variously described in the literature (De Geer, 1940; Zilliacus, 1989; Lundqvist, 2000), but in fjords on Svalbard similar ridges have always been interpreted as either annual push moraines deposited during overall glacier retreat or as crevasse-squeeze ridges formed after surge termination (Solheim and Pfirman, 1985; Boulton et al, 1996; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008; Flink et al, 2015). Based on their dimensions and morphology, some of the De Geer moraines in Kongsfjorden appear similar to annual push moraines described from e.g. Borebukta and Yoldiabukta in Spitsbergen (Ottesen and Dowdeswell, 2006). This is the case especially for the areas in front of Blomstrandbreen, Conwaybreen, and Kongsbreen South, where the longer, more continuous ridges could have been deposited from, and reflect the shape of, the grounding line (Fig. 5b,d). However, the number of De Geer moraines in front of the different glaciers is inconsistent with the number of elapsed years since the formation of the outermost moraine. Assuming that glacier retreat began shortly after moraine formation (cf. Flink et al, 2015), the 55 ridges in front of Blomstrandbreen suggest the outermost moraine (RK1) to be formed in 1955. However, RK1 is probably considerably older, as the glacier front was at least 1.5 km further inland in 1956 (see also section 6.2 below). Furthermore, we would expect c. 113 ridges in front of Conwaybreen (rather than the observed 45), and c. 46 (rather than the observed 30) ridges in front of Kongsbreen South. We thus conclude that if the De Geer moraines in Kongsfjorden were formed from the same mechanism as (annual) push moraines from other fjords, the ridges in Kongsfjorden were formed at more irregular timescales (i.e. not annually).

In places the Kongsfjorden ridges exhibit a saw-tooth pattern in planform, which could be related to formation by a combination of longitudinal crevasse infill and sediment pushing at the glacier snout (Sharp, 1984; Evans and Twigg, 2002; Evans and Orton, 2014; Evans et al, in press). This could suggest that both processes were active in Kongsfjorden, and that the De Geer moraines thus reflect a combination of recessional push moraines and crevasse-squeeze ridges.

Although both, formation as end moraines or as crevasse-squeeze ridges, is possible, the almost perfect symmetry between proximal and distal slopes, the occasional cross-cutting of the ridges, a generally discontinuous character, and a largely similar pattern between crevasses and De Geer moraines (Fig. 9) suggest most of the moraines in Kongsfjorden to derive from crevasse-squeezing. Moreover, on the bathymetry data, the majority of the De Geer moraines appear more consistent with crevasse-squeeze ridges (rather than annual retreat moraines) from Van Keulenfjorden, Rindersbukta (Ottesen et al, 2008) and Tempelfjorden (Flink et al, 2015). In planform their pattern of distribution also compares with crevasse-squeeze ridges from terrestrial surge-type glaciers (Evans and Rea, 1999, 2003), thus further supporting a crevasse-squeeze origin.

Based on the above observations, we therefore suggest that the majority of the De Geer moraines in Kongsfjorden resemble crevasse-squeeze ridges. Nonetheless there are also elements which are compatible with formation of some of the ridges as retreat moraines. This indicates that De Geer moraines

associated with surging glaciers in Spitsbergen fjords form by a variety of processes and thus that submarine landform assemblages in front of surge-type glaciers are more diverse than previously described (cf. e.g. Plassen et al, 2004; Ottesen and Dowdeswell, 2006; Ottesen et al, 2008; Flink et al, 2015).

## 6.2 Timing of landform formation

From correlation of the terminal moraines with known ice front positions (Liestøl, 1988), we infer that all the submarine glacial landforms in inner Kongsfjorden were deposited throughout the past 150 years. Registered surge years are 1869 for Kronebreen, 1948 for Kongsvegen and 1960 for Blomstrandbreen (Hagen, 1993). RK1 was deposited around 1869 (Fig. 10) and probably marks the maximum extent of the Kronebreen surge. RK3 fits the 1948 position, when Kongsvegen surged (Fig. 10). RC1 also dates back to 1869, indicating that the Kronebreen surge caused a simultaneous advance of Kongsbreen North. RK2 and RC2 are close to ice front positions from 1897 (Fig. 10). Neither Kongsbreen nor Conwaybreen are documented to be of surge-type (Hagen, 1993), hence it is possible that these moraines were deposited from a climatically induced advance during the Little Ice Age (LIA). However, the LIA is documented to have ended ~1900 AD (cf. Mangerud and Landvik, 2007) and temperatures were probably warmer already in 1897, making a purely climatic advance unlikely. Furthermore, the similarity between the landforms in front of Conwaybreen and Kongsbreen with those in front of the surge-type glaciers (Blomstrandbreen, Kronebreen and Kongsvegen) could suggest that Kongsbreen did surge. The presence of De Geer moraines between RC2 and the present glacier fronts supports this and we thus infer that a Kongsbreen surge is the more likely scenario.

The ice front positions from 1956 and 1966, as well as the fact that Blomstrandbreen advanced ~600 m during its active surge phase (Liestøl, 1988) strongly suggest RB3 to be the terminal surge moraine from 1960 (Fig. 10). Although the exact chronology of deposition of RB1 and RB2 is still pending, the overridden character of RB2 indicates that it is older than RB1.


We conclude that the geomorphological evidence from the landform assemblages in Kongsfjorden is consistent with surging, especially because (1) the positions of the terminal moraines largely coincide with the glacier front positions documented for respective surge years (cf. Liestøl, 1988), (2) the likely presence of De Geer moraines resembling crevasse-squeeze ridges is considered as possibly surge-diagnostic, (3) glacial lineations in front of all the glaciers indicate fast ice flow, a characteristic of the active phase of a surge cycle (Meier and Post, 1969), and (4) overridden moraines in front of Blomstrandbreen and Kronebreen/Kongsvegen are the product of multiple advances.

## 7 Conclusions

Multibeam and seismic investigations in inner Kongsfjorden reveal a variety of landforms that formed as a result of glacier surges during the past c. 150 years. These include (1) overridden (recessional) moraines from previous ice advance(s), (2) glacial lineations of two types: (a) smooth and (b) sharp-crested groove-ridge features of different sizes, both formed when basal sediments are deformed as a consequence of a rapidly advancing glacier. (3) Large transverse ridges mark the maximum glacier extent during a surge event and were deposited at the end of surges of Kronebreen in 1869, Kongsvegen in 1948, and Blomstrandbreen in 1960. Terminal moraines in front of Conwaybreen/Kongsbreen North and Kongsbreen South/Kronebreen/Kongsvegen coincide with glacier front positions from ~1897, and suggest a Kongsbreen surge for that year. The distal flanks of most of the terminal moraines are characterized by the occurrence of (4) lobe-shaped debris flows resulting from sediment failure or pushing of sediments at the glacier front during the late stages of advance or shortly after. The formation of (5) De Geer moraines, occurred following surge termination and is linked to either crevasse-squeezing during the glacier's stagnant transitional phase and/or to sediment push from small re-advances/halts during overall retreat.


Suspension settling from meltwater plumes and ice rafting are dominant sedimentary processes in the fjord, leading to the deposition of stratified glacialine muds with variable numbers of clasts. Reworking of sediments by glacier surging results in the deposition of sediment lobes containing massive silty clay with frequent clasts. Minimum sediment accumulation rates were ~30 cm a<sup>-1</sup> approximately 2.5 km beyond the front of Kongsvegen after it reached its maximum surge extent in 1948, but rapidly decreased to an average rate of 1.8 cm a<sup>-1</sup> around 1950.

**Acknowledgements** This work, as part of the PetroMaks project "Glaciations in the Barents Sea area (GlaciBar)", was funded by the Research Council of Norway (RCN), Statoil, Det Norske oljeselskap ASA and BG group Norway (grant 200672). It also contributes to the Research School in Arctic Marine Geology and Geophysics (AMGG) and to the Centre of Excellence "Arctic Gas Hydrate, Environment and Climate (CAGE)" at the Department of Geology, UiT – The Arctic University of Norway. CAGE is funded by RCN (grant 223259). The project was further funded by the European Commission FP7-People 2012–Initial Training Networks "Glaciated North Atlantic Margins (GLANAM)". Measurements of sediment accumulation rates were supported by grant IP2010040970 from the Polish Ministry of Science and Higher Education and grant 2013/10/E/ST10/00166 from the Polish National Science Center. We thank the captain and crew of R/V Helmer Hanssen (previously Jan Mayen) and Steinar Iversen for collecting and processing the data and are grateful to Monica Winsborrow and Rune Matningsdal for collecting the sediment cores. Denise Christina R  ther and Lilja R  n Bjarnad  ttir's assistance with the use of the software was greatly appreciated and Beata Sternal kindly helped with the laboratory analyses.




**Fig. 1** a) Map of Spitsbergen with red rectangle indicating the extent of b) bathymetry available for this study and surrounding areas. 1=Breøyane, 2=Gerdøya, 3=Løvenøyane. Black dots indicate location of the two sediment cores 10JM-GlaciBar-GC01 (south) and 10JM-GlaciBar-GC02 (north). Satellite imagery downloaded from Svalbardkartet. c) Locations and extent of available chirp lines, with bright green lines showing the location of profiles in Fig. 6. The colour scale indicates water depth and refers to both, b) and c).






**Fig. 2** Distribution of the interpreted landforms in Kongsfjorden. a) Overview of the bathymetry relief and indication of the locations for subsequent figures; b) Map of overridden moraines and glacial lineations; c) Map of terminal moraines and debris lobes. The bathymetric high refers to work from Trusel et al (2010) and Kehrl et al (2011); d) Map of De Geer moraines in the fjord, with indications of locations of the zoom-ins e) and f). Maps in this paper were created using a Svalbard DTM, available online from the Norwegian Polar Institute.




**Fig. 3** The locations of all subfigures are outlined in Fig. 2a. a) Shaded relief image of the bathymetry showing overridden moraines in front of Kronebreen/Kongsvegen, which are overprinted by streamlined bedforms and small transverse ridges. b) Profile A-A' from proximal to distal across overridden moraines. c) Groove-ridge features in front of Blomstrandbreen with the profile C-C' across them shown in d). e) Small streamlined ridges in front of Kongsbreen North. The profile E-E' across them is shown in f).




**Fig. 4** The locations of all subfigures are outlined in Fig. 2a. a) Shaded relief image of the bathymetry showing three terminal moraines in front of Kronebreen/Kongsvegen (RK1, RK2 and RK3), with RK1 marking the most distal moraine. b) Terminal moraines RK1, 2 and 3 west of Kronebreen/Kongsvegen. The large debris lobe associated with RK1 is also visible. c) Debris lobes off Blomstrandbreen's and Conwaybreen's outermost moraines, RB1 and RB2. Water depths range from -15 to -50 m. d) and e) show profiles A-A' and B-B', respectively, across RK1, RK2 and RK3 from proximal to distal.







**Fig. 5** The locations of all subfigures are outlined in Fig. 2a. Shaded relief image of the bathymetry showing the different appearance of De Geer moraines in Kongsfjorden in front of a) Blomstrandbreen, b) Conwaybreen, c) Kronebreen/Kongsvegen and d) southeast of Blomstrandbreen. d) also shows the large moraine ridge RB3. Cross-sectional profiles across the features from proximal to distal are shown in e), f), g) and h) respectively.




**Fig. 6** a) Black polygon showing the extent of the bathymetry with white lines showing the locations of b), c), and d); b) Chirp line 10JM-GlaciBar025 with the Y-axis showing two-way travel time (TWT), and its facies interpretation in b'). Proximal and distal refer to the proximity to the Kronebreen/Kongsvegen ice margin; c) Chirp line 10JM-GlaciBar038 with facies interpretation in c'; d) Chirp line 10JM-GlaciBar055 with facies interpretation in d'). Red rectangle indicates the extent of Fig. 8.



**Fig. 7** a) Logs and colour images of the sediments in the two cores GC01 and GC02. b) Total  $^{210}\text{Pb}$ , excess  $^{210}\text{Pb}$  and  $^{137}\text{Cs}$  activity profiles for the sediment cores. Vertical error bars represent depth range of the sample, whereas horizontal error bars indicate 2-sigma measurement uncertainties. The derived age model provides approximate average sedimentation accumulation rates shown by the different colours in the  $^{137}\text{Cs}$ -plot.




**Fig. 8** Zoom-in of seismic line 10JM-GlaciBar055 showing location and penetration of the sedimentary units within the GC01 and GC02. Units 1, 2 and 3 correlate with the seismic facies 1, 3 and 2, respectively.



**Fig. 9** a) Map of De Geer moraines in Kongsfjorden with rectangles showing locations of b), c) and d). B), c), and d) show maps of De Geer moraines in front of (black lines) and crevasses on top of the glaciers (red lines).





**Fig. 10** Terminal moraines in Kongsfjorden with respect to glacier front positions reconstructed by Liestøl (1988) in front of a) Blomstrandbreen (RB1, 2, 3) and Conwaybreen/Kongsbreen North (RC1, 2), and b) Kronebreen/Kongsvegen (RK1, 2, 3). Max refers to the unknown year of Blomstrandbreen's maximum extent (cf. Liestøl, 1988)

## References

- Andreassen K, Nilssen E, Ødegaard C (2007) Analysis of shallow gas and fluid migration within the Plio-Pleistocene sedimentary succession of the SW Barents Sea continental margin using 3D seismic data. *Geo-Marine Letters* 27(2-4):155–171
- Appleby P (2008) Three decades of dating recent sediments by fallout radionuclides: a review. *Holocene* 18:83–93
- Baeten N, Forwick M, Vogt C, Vorren T (2010) Late Weichselian and Holocene sedimentary environments and glacial activity in Billefjorden, Svalbard. *Geological Society, London, Special Publications* 344(1):207–223
- Beaudry LM, Prichonnet G (1991) Late Glacial De Geer moraines with glaciofluvial sediment in the Chapais area, Québec (Canada). *Boreas* 20(4):377–394
- Benn D, Evans D (2010) *Glaciers and Glaciation* (2 ed.). Hodder Education
- Björnsson H, Pálsson F, Sigurðsson O, Flowers G (2003) Surges of glaciers in Iceland. *Annals of Glaciology* 36(1):82–90
- Blake KP (2000) Common origin for De Geer moraines of variable composition in Raudvassdalen, northern Norway. *Journal of Quaternary Science* 15(6):633–644
- Błaszczak M, Jania JA, Hagen JO (2009) Tidewater glaciers of Svalbard: Recent changes and estimates of calving fluxes. *Polish Polar Research* 30(2):85–142
- Boulton G (1986) Push-Moraines and glacier-contact fans in marine and terrestrial environments. *Sedimentology* 33(5):667–698
- Boulton G, Van Der Meer J, Hart J, Beets D, Ruegg G, Van Der Wateren F, Jarvis J (1996) Till and moraine emplacement in a deforming bed surge – an example from a marine environment. *Quaternary Science Reviews* 15(10):961–987
- Clark C (1993) Mega-scale glacial lineations and cross-cutting ice-flow landforms. *Earth Surface Processes and Landforms* 18(1):1–29
- Clark CD (1994) Large-scale ice-moulding: a discussion of genesis and glaciological significance. *Sedimentary Geology* 91(1):253–268
- Cowan EA, Powell RD (1991) Ice-proximal sediment accumulation rates in a temperate glacial fjord, southeastern Alaska. *Geological Society of America Special Papers* 261:61–74
- De Geer G (1940) *Geochronologia suecica principes*, vol 18. Almqvist & Wiksells
- Dowdeswell J, Hamilton G, Hagen J (1991) The duration of the active phase of surge-type glaciers: contrasts between Svalbard and other regions. *Journal of Glaciology* 37(127):388–400
- Dowdeswell J, Hodgkins R, Nuttall AM, Hagen J, Hamilton G (1995) Mass balance change as a control on the frequency and occurrence of glacier surges in Svalbard, Norwegian High Arctic. *Geophysical Research Letters* 22(21):2909–2912

- Elverhøi A, Liestøl O, Nagy J (1980) Glacial erosion, sedimentation and micro-fauna in the inner part of Kongsfjorden, Spitsbergen. *Norsk Polarinstitutt Skrifter* 172:33–58
- Elverhøi A, Lønne Ø, Seland R (1983) Glaciomarine sedimentation in a modern fjord environment, Spitsbergen. *Polar Research* 1(2):127–150
- Elverhøi A, Andersen E, Dokken T, Hebbeln D, Spielhagen R, Svendsen J, Forsberg C (1995) The growth and decay of the Late Weichselian ice sheet in western Svalbard and adjacent areas based on provenance studies of marine sediments. *Quaternary Research* 44(3):303–316
- Evans D (2003) Ice-marginal terrestrial landsystems: active temperate glacier margins. *Glacial Landsystems* Arnold, London pp 12–43
- Evans D, Orton C (2014) Heinabergsjökull and Skálafellsjökull, Iceland: active temperate piedmont lobe and outwash head glacial landsystem. *Journal of Maps* 11(3):415–431
- Evans D, Rea B (1999) Geomorphology and sedimentology of surging glaciers: a land-systems approach. *Annals of Glaciology* 28(1):75–82
- Evans D, Rea B (2003) Surging glacier landsystem. *Glacial Landsystems* Arnold, London pp 259–288
- Evans D, Twigg D (2002) The active temperate glacial landsystem: a model based on Breiðamerkurjökull and Fjallsjökull, Iceland. *Quaternary Science Reviews* 21(20):2143–2177
- Evans D, Lemmen DR, Rea B (1999) Glacial landsystems of the southwest Laurentide Ice Sheet: modern Icelandic analogues. *Journal of Quaternary Science* 14(7):673–691
- Evans D, Ewertowski M, Orton C (in press) Fláajökull (north lobe), Iceland: active temperate piedmont lobe glacial landsystem. *Journal of Maps*
- Evans DJA, Phillips E, Hiemstra J, Auton C (2006) Subglacial till: formation, sedimentary characteristics and classification. *Earth-Science Reviews* 78(1):115–176
- Farrow GE, Syvitski JP, Tunncliffe V (1983) Suspended particulate loading on the macrobenthos in a highly turbid fjord: Knight Inlet, British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* 40(S1):s273–s288
- Flink AE, Noormets R, Kirchner N, Benn DI, Luckman A, Lovell H (2015) The evolution of a submarine landform record following recent and multiple surges of Tunabreen glacier, Svalbard. *Quaternary Science Reviews* 108:37–50
- Forwick M, Vorren T (2007) Holocene mass-transport activity and climate in outer Isfjorden, Spitsbergen: marine and subsurface evidence. *The Holocene* 17(6):707–716
- Forwick M, Vorren T (2009) Late Weichselian and Holocene sedimentary environments and ice rafting in Isfjorden, Spitsbergen. *Paleogeography, Paleoclimatology, Paleoecology* 280(1):258–274
- Forwick M, Vorren T (2010) Stratigraphy and deglaciation of the Isfjorden area, Spitsbergen. *Norwegian Journal of Geology* 90:163–179
- Forwick M, Vorren T, Hald M, Korsun S, Roh Y, Vogt C, Yoo K (2010) Spatial and temporal influence of glaciers and rivers on the sedimentary

- environment in Sassenfjorden and Tempelfjorden, Spitsbergen. Geological Society, London, Special Publications 344(1):163–193
- Gilbert R, Nielsen N, Möller H, Desloges J, Rasch M (2002) Glacimarine sedimentation in Kangerdluk (Disko Fjord), West Greenland, in response to a surging glacier. *Marine Geology* 191:1–18
- Hagen J (1993) Glacier atlas of Svalbard and Jan Mayen, vol 129. Norsk Polarinstitutt Middelelser
- Hald M, Dahlgren T, Olsen TE, Lebesbye E (2001) Late Holocene paleoceanography in Van Mijenfjorden, Svalbard. *Polar Research* 20(1):23–35
- Henriksen M, Alexanderson H, Landvik JY, Linge H, Peterson G (2014) Dynamics and retreat of the Late Weichselian Kongsfjorden ice stream, NW Svalbard. *Quaternary Science Reviews* 92:235–245
- Hoppe G (1957) Problems of glacial morphology and the ice age. *Geografiska Annaler* pp 1–18
- Howe J, Moreton S, Morri C, Morris P (2003) Multibeam bathymetry and the depositional environments of Kongsfjorden and Krossfjorden, western Spitsbergen, Svalbard. *Polar Research* 22 (2):301–316
- Ingólfsson O, Landvik J (2013) The Svalbard-Barents Sea ice-sheet – historical, current and future perspectives. *Quaternary Science Reviews* 64:33–60
- Ito H, Kudoh S (1997) Characteristics of water in Kongsfjorden, Svalbard. *Proceedings of the NIPR symposium on polar meteorology and glaciology* 11:211–232
- Jernas P, Klitgaard-Kristensen D, Husum K, Wilson L, Koç N (2013) Palaeoenvironmental changes of the last two millennia on the western and northern Svalbard shelf. *Boreas* 42(1):236–255
- Jessen S, Rasmussen T, Nielsen T, Solheim A (2010) A new Late Weichselian and Holocene marine chronology for the western Svalbard slope 30000–0 cal years BP. *Quaternary Science Reviews* 29(9):1301–1312
- Kamb B (1987) Glacier surge mechanism based on linked cavity configuration of the basal water conduit system. *Journal of Geophysical Research* B 92(B9):9083–9100
- Kehrl L, Hawley R, Powell R, Brigham-Grette J (2011) Glacimarine sedimentation processes at Kronebreen and Kongsvegen, Svalbard. *Journal of Glaciology* 57(205):841–847
- Kempf P, Forwick M, Laberg J, Vorren T (2013) Late Weichselian – Holocene sedimentary palaeoenvironment and glacial activity in the high-Arctic van Keulenfjorden, Spitsbergen. *The Holocene* 23(11):1607–1618
- King E, Hindmarsh R, Stokes C (2009) Formation of mega-scale glacial lineations observed beneath a West Antarctic ice stream. *Nature Geoscience* 2(8):585–588
- Koide M, Soutar A, Goldberg ED (1972) Marine geochronology with 210 Pb. *Earth and Planetary Science Letters* 14(3):442–446
- Kristensen L, Benn D, Holmes A, Ottesen D (2009) Mud aprons in front of Svalbard surge moraines: Evidence of subglacial deforming layers or proglacial glaciotectionics? *Geomorphology* 111(3):206–221

- Landvik J, Bondevik S, Elverhøi A, Fjeldskaar W, Mangerud J, Salvigsen O (1998) The last glacial maximum of Svalbard and the Barents Sea area: ice sheet extent and configuration. *Quaternary Science Reviews* 17(1-3):43–75
- Larsen E, Longva O, Follestad BA (1991) Formation of De Geer moraines and implications for deglaciation dynamics. *Journal of Quaternary Science* 6(4):263–277
- Lehman S, Forman S (1992) Late Weichselian glacier retreat in Kongsfjorden, west Spitsbergen, Svalbard. *Quaternary Research* 37(2):139–154
- Liestøl O (1969) Glacier surges in west Spitsbergen. *Canadian Journal of Earth Sciences* 6:895–898
- Liestøl O (1988) The glaciers in the Kongsfjorden area, Spitsbergen. *Norske geografiske Tidsskrifter* 42:231–238
- Lundqvist J (1981) Moraine morphology. terminological remarks and regional aspects. *Geografiska Annaler Series A Physical Geography* pp 127–138
- Lundqvist J (2000) Palaeoseismicity and De Geer Moraines. *Quaternary International* 68:175–186
- MacLachlan S, Howe J, Vardy M (2010) Morphodynamic evolution of Kongsfjorden-Krossfjorden, Svalbard, during the Late Weichselian and the Holocene. *Geological Society London, Special Publications* 344(1):195–205
- Mangerud J, Landvik J (2007) Younger Dryas cirque glaciers in western Spitsbergen: smaller than during the Little Ice Age. *Boreas* 36(3):278–285
- McKee B, Nittrouer C, DeMaster D (1983) Concepts of sediment deposition and accumulation applied to the continental shelf near the mouth of the Yangtze River. *Geology* 11:631–633
- Meier M, Post A (1969) What are glacier surges? *Canadian Journal of Earth Sciences* 6:807–817
- Murray T, Dowdeswell JA, Drewry DJ, Frearson I (1998) Geometric evolution and ice dynamics during a surge of Bakaninbreen, Svalbard. *Journal of Glaciology* 44(147):263–374
- Ó Cofaigh C, Dowdeswell J, Allen C, Hiemstra J, Pudsey C, Evans J, Evans D (2005) Flow dynamics and till genesis associated with a marine-based Antarctic paleo-ice stream. *Quaternary Science Reviews* 24(5):709–740
- Ottesen D, Dowdeswell J (2006) Assemblages of submarine landforms produced by tidewater glaciers in Svalbard. *Journal of Geophysical Research* 111:F01,016
- Ottesen D, Dowdeswell J, Rise L (2005) Submarine landforms and the reconstruction of fast-flowing ice streams within a large Quaternary ice sheet: the 2500-km-long Norwegian-Svalbard margin (57°– 80°N). *Geological Society of America Bulletin* 117:1033–1050
- Ottesen D, Dowdeswell J, Landvik J, Mienert J (2007) Dynamics of the Late-Weichselian ice sheet on Svalbard inferred from high-resolution sea-floor morphology. *Boreas* 36:286–306
- Ottesen D, Dowdeswell J, Benn D, Kristensen L, Christiansen H, Christensen O, Vorren T (2008) Submarine landforms characteristic of glacier surges in two Spitsbergen fjords. *Quaternary Science Reviews* 27(15):1583–1599

- Plassen L, Vorren T, Forwick M (2004) Integrated acoustic and coring investigations of glacial deposits in Spitsbergen fjords. *Polar Research* 23(1):89–110
- Raymond C (1987) How do glaciers surge? A review. *Journal of Geophysical Research: Solid Earth* (1978–2012) 92(B9):9121–9134
- Robbins J, Edgington D (1975) Determination of recent sedimentation rates in Lake Michigan using Pb-210 and Cs-137. *Geochimica et Cosmochimica Acta* 39:285–304
- Sharp M (1984) Annual moraine ridges at Skálafellsjökull, south-east Iceland. *Journal of Glaciology* 30:104
- Sharp M (1985) Crevasse-fill-ridges - a landform type characteristic of surging glaciers? *Geografiska Annaler Series A Physical Geography* 67(3/4):213–220
- Sharp M (1988) Surging glaciers: behaviour and mechanisms. *Progress in Physical Geography* 12(3):349–370
- Solheim A (1991) The depositional environment of surging sub-polar tidewater glaciers. *Norsk Polarinstitutt Skrifter* 194:1–97
- Solheim A, Pfirman S (1985) Sea-floor morphology outside a grounded, surging glacier; Bråsvellbreen, Svalbard. *Marine Geology* 65:127–143
- Sollid JL (1989) Comments on the genesis of De Geer moraines. *Norsk Geografisk Tidsskrift-Norwegian Journal of Geography* 43(1):45–47
- Stokes C, Clark C (2002) Are long subglacial bedforms indicative of fast ice flow? *Boreas* 31(3):239–249
- Strömberg B (1965) Mappings and geochronological investigations in some moraine areas of south-central Sweden. *Geografiska Annaler Series A, Physical Geography* 47(2):73–82
- Svendsen H, Beszczynska-Møller A, Hagen J, Lefauconnier B, Tverberg V, Gerland Sea (2002) The physical environment of Kongsfjorden-Krossfjorden, an Arctic fjord system in Svalbard. *Polar Research* 21(1):133–166
- Syvitski JP (1989) On the deposition of sediment within glacier-influenced fjords: oceanographic controls. *Marine Geology* 85(2):301–329
- Szczuciński W, Zajaczkowski M (2012) Factors controlling downward fluxes of particulate matter in glacier-contact and non-glacier contact settings in a subpolar fjord (Billefjorden, Svalbard). *International Association of Sedimentology, Special Publication* 44:369–386
- Szczuciński W, Zajaczkowski M, Scholten J (2009) Sediment accumulation rates in subpolar fjords – impact of post-Little Ice Age glacier retreat, Billefjorden, Svalbard. *Estuarine, Coastal and Shelf Science* 85:345–356
- Trusel L, Powell R, Cumpston R, Brigham-Grette J (2010) Modern glacimarine processes and potential future behaviour of Kronebreen and Kongsvegen polythermal tidewater glaciers, Kongsfjorden, Svalbard. *Geological Society of London Special Publications* 344:89–102
- Tulaczyk S, Scherer R, Clark C (2001) A ploughing model for the origin of weak tills beneath ice streams: a qualitative treatment. *Quaternary International* 86:59–70
- Zaborska A, Pempkowiak J, Papucci C (2006) Some sediment characteristics and sedimentation rates in an Arctic fjord (Kongsfjorden, Svalbard). *Annual*

Environmental Protection 8:79–96

Zajaczkowski M, Szczuciński W, Bojanowski R (2004) Recent changes in sediment accumulation rates in Adventfjorden, Svalbard. *Oceanologia* 46:217–231

Zilliacus H (1989) Genesis of De Geer moraines in Finland. *Sedimentary geology* 62(2):309–317